

University of Groningen

## Nonuniqueness of the Phase Shift in Central Scattering due to Monodromy

Dullin, Holger R.; Waalkens, Holger

*Published in:*  
Physical Review Letters

*DOI:*  
[10.1103/PhysRevLett.101.070405](https://doi.org/10.1103/PhysRevLett.101.070405)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2008

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Dullin, H. R., & Waalkens, H. (2008). Nonuniqueness of the Phase Shift in Central Scattering due to Monodromy. *Physical Review Letters*, 101(7), [070405]. <https://doi.org/10.1103/PhysRevLett.101.070405>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

**Dullin and Waalkens Reply:** The results of our Letter [1] are correct. In the Letter we showed that the phase shift for 2D central scattering at a smooth repulsive potential cannot be uniquely defined as a globally *smooth* function of angular momentum and energy due to a topological obstruction similar to monodromy in bound systems.

The preceding Comment by Eiglsperger *et al.* [2] raises two points. First of all, the authors of the Comment point out that the phase shift can be uniquely defined as a *nonsmooth* function. Second, the authors claim that the WKB approximation is bad for the class of potentials studied in our Letter. The first point is trivial and no contradiction to what we say in our Letter. We suppose that the commentators raise this point because we did not mention the word “smooth” in the title and abstract of our Letter. We stress, however, that the entire discussion in our Letter is based on and is only meaningful for globally *smooth phase shifts*. The smoothness assumption is explicitly stated in the paragraph before Eq. (3) in our Letter, and also implied by the term *monodromy* in the title. In the second point the authors give an example of a potential for which the WKB approximation for *s*-wave scattering is bad near the critical threshold energy  $E = 0$ . It is well known the WKB approximations become bad near critical values. However, it is the essence of monodromy that, for the smooth continuation of the phase shift, one can stay away from critical values. Accordingly, we will show below that for the relevant noncritical energies, the WKB approximation works very well also for the “counterexample” in the Comment [2].

**First point.**—The key point of our Letter is that globally  $\Delta W$  cannot be defined as a single-valued *smooth* function of  $l$  and  $p = (2mE)^{1/2}$ . This is surprising, because *locally*  $\Delta W$  can be defined as such a single-valued smooth function. Two-dimensional central scattering is integrable. The angular momentum  $L = xp_y - yp_x$  and the Hamiltonian  $H = \frac{1}{2m}(p_x^2 + p_y^2) + V(x^2 + y^2)$  are constants of motion in involution with linearly independent gradients almost everywhere in the phase space  $\mathbb{R}^4$ . In fact, for a smooth repulsive central potential of the type discussed in our Letter, the only points in the phase space at which  $\nabla L$  and  $\nabla H$  are linearly dependent are the origin, with corresponding critical value  $(L, H) = (0, E_c)$  where  $E_c = V(0)$ , and the trivial equilibrium points at infinity where the particle is at rest with critical value  $E = 0$ . In a neighborhood of a regular value  $(l, E)$  the Liouville-Arnold theorem guarantees that  $\Delta W$  is a smooth function of  $l$  and  $E$ . However, these *local* definitions of  $\Delta W$  cannot be patched together to give a *global* single-valued smooth function  $\Delta W$ , i.e., a function which is smooth and single-valued on the set of regular values of  $L$  and  $H$ . The function  $\Delta W$  near




FIG. 1. (a) Exact phase shift and the corresponding WKB approximation for the potential  $V(r) = 20 \exp(-r^2)$ . ( $m = 1$ ,  $\hbar = 0.25$ .) (b) Error of the WKB phase shift.

$(L, H) = (0, E_c)$  has properties that are similar to the function  $\Re(z \log z - z)$  with  $z = (H - E_c) + iL \in \mathbb{C}$  near zero. Analytic continuation leads to a smooth but multi-valued function. Of course, if one gives up the smoothness of  $\Delta W$  and introduces a branch cut as done in the Comment [2], then  $\Delta W$  becomes unique. This trivial statement, however, has no bearing whatsoever on the result of our Letter.

**Second point.**—In Fig. 1 we compare the (exact) phase shift  $\delta$  with its WKB approximation  $\Delta W$  for *s*-waves scattering ( $l = 0$ ) at the potential  $V(r) = 20 \exp(-r^2)$  considered in the Comment [2]. For energies very close to the critical threshold  $E = 0$  the WKB approximation is bad as pointed out in the Comment [2]. This behavior is related to the fast decay of the potential at infinity. In the regular energy range relevant for the analytic continuation around the critical value  $(L, H) = (0, E_c) = (0, 20)$ , however, the error of the WKB approximation is far below the stated 1% in our Letter.

Holger R. Dullin

School of Mathematics and Statistics  
University of Sydney  
New South Wales 2006, Australia

Holger Waalkens

Department of Mathematics  
University of Groningen  
9747 AG Groningen, The Netherlands  
and School of Mathematics  
University of Bristol  
Bristol BS8 1TW, United Kingdom

Received 23 September 2008; published 7 May 2009

DOI: [10.1103/PhysRevLett.102.188902](https://doi.org/10.1103/PhysRevLett.102.188902)

PACS numbers: 03.65.Nk, 03.65.Sq, 05.45.-a, 45.05.+x

- [1] H.R. Dullin and H. Waalkens, Phys. Rev. Lett. **101**, 070405 (2008).
- [2] J. Eiglsperger, H. Friedrich, and J. Madroñero, preceding Comment, Phys. Rev. Lett. **102**, 188901 (2009).